



NAME: _____

We perform procedures at the following locations:

- | | |
|--|--|
| <input type="checkbox"/> GIEA
15005 Shady Grove Road. Ste 200
Rockville, MD. 20850
(301) 340-8099 | <input type="checkbox"/> Shady Grove Adventist Hospital
Out- Patient Surgery Center
9901 Medical Center Drive
Rockville, MD 20850 |
|--|--|

Please ARRIVE at: _____ on _____
TIME DATE

PREPARATION INSTRUCTIONS FOR COLONOSCOPY (STANDARD MIRALAX METHOD)

PURCHASE 2 OR MORE DAYS PRIOR TO PROCEDURE (OVER-THE-COUNTER):

- One (1) bottle of Miralax 8.3 ounces (equivalent to 238gms).
- Eight (8) Dulcolax or Bisacodyl 5mg tablets (they are both equivalent).

7 DAYS BEFORE YOUR PROCEDURE:

- Discontinue medications containing iron; follow your providers instructions regarding aspirin, Plavix or anticoagulation medicines (e.g. Coumadin, Pradaxa, Xeloda)

THE ENTIRE DAY PRIOR TO YOUR PROCEDURE YOU MUST BE ON A CLEAR LIQUID DIET – NO SOLID FOODS.

- Drink 6-8 glasses of clear liquids and half **MUST** contain sugar and salt.
- Coffee or Tea (no milk)
- Clear broth or bouillon – **MUST** drink 16 oz of either broth or bouillon
- Carbonated or non-carbonated soft drinks (Coke, Pepsi, Sprite, etc)
- Clear fruit juices (i.e. Apple Juice, Cranberry, White grape juice, Gatorade(**NOT G2**))
- Jell-O, popsicles, hard candy

NOTE: DARK COLOR CLEAR LIQUIDS SUCH AS RED OR PURPLE ARE OK TO HAVE JUST BE SURE TO DRINK LOTS OF WATER AS WELL.

THE DAY BEFORE PROCEDURE:

- **Between 12:00p.m. And 1:00p.m.** -Take 2 Dulcolax/Bisacodyl tablets with 8oz. of a clear liquid.
- **Between 3:00p.m. And 4:00p.m.** – Take another 2 Dulcolax/bisacodyl tablets with 8oz. of a clear liquid.
- **Anytime between 4:00p.m. And 6:30p.m.** Mix the 1 bottle of Miralax(or generic) with your 64 oz of Gatorade. **MUST BE GATORADE ONLY.** Then begin to drink 1 8 oz glass of the solution every 20-30 mins until the solution is gone. Continue to drink clear liquids until going to bed.
- **Between 6:00p.m. – 7:00p.m.** – Take another 2 Dulcolax/bisacodyl tablets with 8oz of a clear liquid.
- **Between 10:00p.m. - 11:00p.m.** -Take another 2 Dulcolax/Bisacodyl tablets with 8oz. of a clear liquid.

Capital Digestive Care

Alan N. Schulman, MD
Sheila G. Levin, MD
Julia C. Korenman, MD
Lawrence A. Bassin, MD
David L. Jager, MD
Olanma Y. Okoji, MD
Colleen M. Kennedy-Smith, CRNP
Lisa Rainsford, PA-C

ROCKVILLE, MD
14955 Shady Grove Rd
Suite 150
Rockville, MD 20850

301.340.3252 **PHONE**
301.340.1423 **FAX**

capitaldigestivecare.com

OVER →→→

NOTE: If you have a noon or later procedure you should split the Miralax. You should drink 2/3 between 4 p.m. and 11:00 p.m. the evening before, and drink the remaining 1/3 of Miralax the morning of the procedure between 6 a.m. and 9 a.m.

DAY OF YOUR EXAM: Take all of your morning medications with small sips of water **ONLY UP UNTIL 4 HOURS PRIOR TO YOUR PROCEDURE TIME.**

PLEASE NOTE: IF YOU FIND IT NECESSARY TO CANCEL YOUR PROCEDURE **WITHIN 48 HOURS PRIOR TO THE SCHEDULED APPOINTMENT A \$150 CANCELLATION FEE WILL BE CHARGED.**