

Maryland Digestive Disease Center

Jeffrey S. Garbis, MD, FACG Richard M. Chasen, MD. FACG Jeffrey Bernstein, MD, FACG Theodore Y Kim MD FACG Marvin E. Lawrence II. MD. FACG Sean M. Karp, MD, FACG Priti Bijpuria, MD

ELLICOTT CITY, MD 4801 Dorsey Hall Drive Suite 120

Ellicott City, MD 21042 410.730.9363 **OFFICE** 410.730.2084 FAX

LAUREL, MD

7350 Van Dusen Road Suites 210 & 230 Laurel, MD 20707 301.498.5500 **OFFICE** 301.498.7346 FAX

TAKOMA PARK, MD 7610 Carroll Avenue

Suite 250 Takoma Park, MD 20912 301.270.3640 OFFICE 301.270.3645 FAX

Patient Instructions for Infrared Coagulation of Hemorrhoids (IRC)

Patient Name:		
Date:	Arrival Time for Procedure:	
With Doctor:		
Location:		

YOU MUST PURCHASE:

(2) 4.5 fl. oz. size enema.

Infrared Coagulation of Hemorrhoids

Infrared coagulation is a procedure used for the treatment of hemorrhoids. The procedure involves the application of infrared light to the base of the hemorrhoid, which clots the hemorrhoid. There are usually three different areas inside the rectum where the hemorrhoids occur, and they are referred to as complexes. One area, or complex, is treated at each office visit. Although the patient will occasionally feel a little warmth, there generally is minimal pain or discomfort. The patient may return to work the same day or the next day. Occasionally a little bleeding will occur between the fourth and tenth day after treatment. The patient returns in approximately 1 month for follow-up treatment.

In order to be properly prepared for your procedure, please review the Pre-Procedure requirements on the attached sheet **one week** prior to your exam and follow the instructions carefully.

Should you have any questions please contact one of our office locations listed below.

Appointment Day M Tu W Th F	Date	Capital Digestive Care, LLC Maryland Digestive Disease Center
5 Days Before Procedure	• Purchase (2) 4.5 fl. oz size enemas	
Day of Procedure	 An hour & a half before your appointment take the first enema, hold it in for 5 minutes and then release into the toilet. Repeat immediately with the second enema. Do not eat anything after you have taken the 2 enemas Follow the instructions provided on the enema package for the proper technique for the insertion of the enema into the rectum. If you are unable to read or communicate in the English language, please bring an interpreter with you to translate. 	

Please note there is a cancellation fee for appointments not cancelled within 48 hours.