

# PLENVU PREP KIT INSTRUCTIONS FOR COLONOSCOPY OR COLONOSCOPY / EGD

PATIEN	TT:	A	RRIVE AT:	:	ON	
FOR A P	PROCEDURE APPOINTMENT WIT	TH DR				
$\sqrt{5}$ DAYS B	EFORE THE PROCEDURE:					
<ul> <li>Pui</li> <li>Scl</li> <li>(ov</li> <li>Pat</li> <li>A c</li> </ul>	rease read through this entire instruction rechase the prepping solution prescription hedule a ride home. Every patient must over 18 years old). The escort should arritionest scheduled after 3:00 pm must have cancellation fee of \$ 150.00 may be chastore the procedure appointment.	n from your Pharma be escorted home be we to the facility on the an escort on site b	y a personal e (1) hour af by 4:30 pm.	ter the proced	dure appointment time.	
√ MEDICAT	ΓIONS					
<ul><li>As dis vis</li><li>Spe pro</li></ul>	o not take iron supplements for five (5) of pirin, diabetic medications/insulin, anti- ccussed with your Physician, Nurse Practit.  ecified medications should be taken the ocedure start time with only a small sip on to stop any medications unless instructions.	coagulation medica etitioner or Physicia morning of the pro of water.	tions, heart r n Assistant a	at the time of	the pre-procedure offic	ee
$\sqrt{3}$ DAYS B	EFORE THE PROCEDURE:					
	ljust your diet to prepare for the procedurn/popcorn, raw fruits, raw vegetables a	1 0		•	_	s,
$\sqrt{\text{THE DAY}}$	BEFORE THE PROCEDURE:					
	is procedure requires that you follow the termine which diet you should follow.	e following diet(s)	on the day be	efore the proc	edure. Your doctor wil	1
	☐ Clear Liquid Diet	Breakfast	Lunch	Dinner		
	Low Residue Diet	Breakfast	Lunch	Dinner- be	fore 4pm	

**ALLOWED ON A CLEAR LIQUID DIET:** apple juice, white grape and white cranberry juice, broth, tea, coffee (without milk or creamer), Jell-O, Italian Ices, popsicles, sodas, Kool-Aid, Gatorade, and water. You should continue these clear liquids up until 4 hours prior to your procedure.

**NOT ALLOWED ON A CLEAR LIQUID DIET:** red, orange, and purple colored liquids, as well as the following products, milk, cream, soy milk equivalent, artificial creamer, juices containing pulp (i.e. orange, grapefruit, pineapple, tomato & V-8 juice). Solid foods are not allowed on a clear liquid diet.

**ALLOWED ON A LOW RESIDUE DIET:** Clear liquids as instructed, and you may also have Ensure, Boost, eggs and plain or flavored yogurt without seeds or nuts. <u>NO other foods are allowed on this diet.</u>

#### PLENVU PREP KIT – DO NOT FOLLOW THE INSTRUCTIONS ON THE BOX

## $\sqrt{\text{ON THE DAY PRIOR TO THE PROCEDURE, FOLLOW THE TABLE BELOW:}}$

8:00 am the day before	Follow a clear liquid diet. It is important to drink as many clear liquids throughout the day to avoid			
the procedure dehydration during the evening prep.				
First Dose	Use the mixing container to mix the contents of the Dose 1 pouch (mango flavor) with at least 16 oz.			
	of water by shaking or using a spoon until it is completely dissolved. Mixing may take up to 2-3			
4:00 pm on the day	minutes.			
before the procedure				
_	Then drink the solution over the next 30 minutes, take your time.			
	Refill the container with at least 16 oz of clear liquids again take your time and slowly finish.			

## $\sqrt{\text{SIX}}$ (6) HOURS PRIOR TO THE PROCEDURE, FOLLOW THE TABLE BELOW:

Second Dose Six (6) hours prior to your scheduled procedure time	Use the mixing container to mix the contents of the Dose 2 pouch A and pouch B- (fruit punch flavor) with at least 16 oz. of water by shaking or using a spoon until it is completely dissolved. Mixing may take up to 2-3 minutes.  Then drink the solution over the next 30 minutes, take your time.
Five (5) hours before your scheduled procedure time	Then drink at least 16 oz. of clear liquid over the next hour.  You must be finished drinking 4 hours prior to your scheduled procedure time.
<b>Additional Instructions:</b>	

### $\sqrt{\text{WHAT TO EXPECT}}$ :

- Expect to have frequent bowel movements and diarrhea within 1-4 hours. Be patient and try walking around to stimulate bowel motility. Remain near a restroom. Continue your prep and if after 4 hours you have not had a bowel movement, contact the office.
- Expect to pass clear or yellow fluid at the completion of your prep.
- You must also take the second dose of the prep even if you are passing clear or yellow fluid.
- Expect that patients who do not fully complete their prep and are not adequately prepared may be canceled or rescheduled.

## $\sqrt{\text{WHAT TO DO TO MAKE PREPPING EASIER:}}$

- Patients can use Destin, Anusol, Tucks pads or Vaseline to coat the rectal area to avoid irritation during the prep.
- Patients can refrigerate the prep to make drinking the solution easier.
- Patients can add a lemon Crystal Lite flavor packet to the solution to help with the taste.
- Patients can drink the solution with a straw or hold your nose when drinking patients should slow down and take time between doses if feeling nauseous.
- Patients should continue drinking clear liquids up to four hours before the procedure. During the 4 hours before
  your procedure, take nothing by mouth, not even water, chewing gum or candy. If you do, the procedure will be
  canceled.

### $\sqrt{\text{LAST MINUTE CHECK LIST:}}$

- Take your blood pressure, heart, Parkinson's, seizure or asthma medications, please take these 3 or more hours prior to the procedure start time with only a small sip of water.
- Other than the medications above, do not take anything by mouth for 4 hours prior to the procedure as listed above.
- If a physician has prescribed for you an inhaler for asthma, bring it with you to the procedure.
- Bring with you your photo ID, insurance Cards, co-payment if required.
- Leave valuables and jewelry at home and wear comfortable clothing.
- Avoid wearing perfume or lotions on the day of your procedure.

PLEASE LIST:		
NAME OF ESCORT HOME:	RELATIONSHIP:	
PHONE NUMBER:		7 2019