

OSMO PREP KIT INSTRUCTIONS FOR COLONOSCOPY OR COLONOSCOPY / EGD

PATIENT:		A	ARRIVE AT	:	ON	
FC	OR A PROCEDURE APPOINTMENT WITH	DR				
√ 5 DAY	YS BEFORE THE PROCEDURE:					
•	Please read through this entire instruction so Purchase the prep prescription from your pharm Schedule a ride home. Every patient must be estable 18 years old). The escort should arrive to the far Patients scheduled after 3:00 pm must have an A cancellation fee of \$ 150.00 may be charged before the procedure appointment.	macy. scorted home by acility one (1) h escort on site b	our after the y 4:30 pm.	procedure appo	pintment time.	
$\sqrt{\mathbf{MED}}$	ICATIONS					
•	Aspirin, diabetic medications/insulin, anticoagulation medications, heart medications, blood thinners; must be discussed with your Physician, Nurse Practitioner or Physician Assistant at the time of the pre-procedure office visit. Specified medications should be taken the morning of the procedure, as prescribed, 3 or more hours prior to the procedure start time with only a small sip of water.					
$\sqrt{3}$ DAY	YS BEFORE THE PROCEDURE:					
•	Adjust your diet to prepare for the procedure. Stop eating high fiber foods including, seeds, quinoa, nuts, beans, corn/popcorn, raw fruits, raw vegetables and bran for three (3) days prior to the procedure.					
√ THE D	AY BEFORE THE PROCEDURE:					
•	• This procedure requires that you follow the following diet(s) on the day before the procedure. Your doctor will determine which diet you should follow.					
	Clear Liquid Diet	Breakfast	Lunch	Dinner		

ALLOWED ON A CLEAR LIQUID DIET: apple juice, white grape and white cranberry juice, broth, tea, coffee (without milk or creamer), Jell-O, Italian Ices, popsicles, sodas, Kool-Aid, Gatorade, and water. You should continue these clear liquids up until 4 hours prior to your procedure.

Breakfast

Lunch

Dinner- before 4pm

NOT ALLOWED ON A CLEAR LIQUID DIET: red, orange, and purple colored liquids, as well as the following products, milk, cream, soy milk equivalent, artificial creamer, juices containing pulp (i.e. orange, grapefruit, pineapple, tomato & V-8 juice). Solid foods are not allowed on a clear liquid diet.

Low Residue Diet

ALLOWED ON A LOW RESIDUE DIET: Clear liquids as instructed, and you may also have Ensure, Boost, eggs and plain or flavored yogurt without seeds or nuts. <u>NO other foods are allowed on this diet.</u>

OSMO PREP KIT - DO NOT FOLLOW THE INSTRUCTIONS ON THE PRESCRIPTION

FOLLOW THE TABLE BELOW THE DAY PRIOR TO THE PROCEDURE:					
8:00 am the day before the procedure	It is important to drink as many clear liquids throughout the day to avoid				
· ·	dehydration during the evening prep.				
First Dose	Take 4 pills with 8 oz. of clear liquid every 15 min for one (1) hour.				
6:00 pm on the day before the procedure	Check off the boxes below to help you keep track of the doses taken.				
	You will take 20 pills with 40 oz. of clear liquid.				
	4 pills/8 oz. 4 pills/8 oz. 4 pills/8 oz. 4 pills/8 oz. 4 pills/8 oz				
	Then drink at least two (2) 16 oz. glasses of water or clear liquid.				
	•				
$\sqrt{\text{FOLLOW THE TABLE BELOW SIX (6) HOURS PRIOR TO THE PROCEDURE:}}$					
Second Dose	Take 4 pills with 8 oz. of clear liquid every 15 min for 30 minutes.				
Six (6) hours before your scheduled	Check off the boxes below to help you keep track of the doses taken.				
procedure time	You will take the remaining 12 pills with 24 oz. of clear liquid				
•	☐ 4 pills / 8 oz. ☐ 4 pills / 8 oz. ☐ 4 pills / 8 oz.				
Five (5) hours before your scheduled	Drink at least two (2) 16 oz. glasses of water or clear liquid over the next				
procedure time	hour. You must be finished drinking 4 hours prior to your scheduled				
procedure time	procedure time.				
Additional Instructions:					
$\sqrt{\text{WHAT TO EXPECT:}}$,				

- Expect to have frequent bowel movements and diarrhea within 1-4 hours. Be patient and try walking around to stimulate bowel motility. Remain near a restroom. Continue your prep and if after 4 hours you have not had a bowel movement, contact the office.
- Expect to pass clear or yellow fluid at the completion of your prep.
- You must also take the second dose of the prep even if you are passing clear or yellow fluid.
- Expect that patients who do not fully complete their prep and are not adequately prepared may be canceled or rescheduled.

$\sqrt{\text{WHAT TO DO TO MAKE PREPPING EASIER:}}$

- Patients can use Desitin, Anusol, Tucks pads or Vaseline to coat the rectal area to avoid irritation during the prep.
- Patients can refrigerate the prep to make drinking the solution easier.
- Patients can drink the solution with Sprite, Ginger Ale or Lemonade.
- Patients can drink with a straw or hold your nose when drinking.
- Patients should slow down and take time between doses if feeling nauseous.
- Patients should continue drinking clear liquids up to four hours before the procedure. During the 4 hours before your procedure, take nothing by mouth, not even water, chewing gum or candy. **If you do, the procedure will be canceled.**

$\sqrt{\text{LAST MINUTE CHECK LIST:}}$

- Take your blood pressure, Parkinson's, seizure or asthma medications. Please take these medications 3 or more hours prior to the procedure start time with only a small sip of water.
- Other than the medications above, do not take anything by mouth for 4 hours prior to the procedure.
- If a physician has prescribed for you an inhaler for asthma, bring it with you to the procedure.
- Bring with you your photo ID, insurance cards, co-payment if required.
- Leave valuables and jewelry at home and wear comfortable clothing.
- Avoid wearing perfume or lotions on the day of your procedure.

PLEASE LIST:

LEASE EIST.	
NAME OF ESCORT HOME:	RELATIONSHIP:
PHONE NUMBER:	