



SUPREP PREP INSTRUCTIONS FOR COLONOSCOPY OR COLONOSCOPY/ EGD

PATIENT: _____ **ARRIVE AT:** _____ **ON** _____

FOR A PROCEDURE APPOINTMENT WITH DR. _____

√ 5 DAYS BEFORE THE PROCEDURE:

- **Please read through this entire instruction sheet.**
- Pick up the prepping solution prescription from the pharmacy.
- Schedule a ride home. Every patient must be escorted home by a personal friend, family member or caregiver (over 18 years old). The escort should arrive to the facility one (1) hour after the procedure appointment time.
- Patients scheduled after 3:00 pm must have an escort on site by 4:30 pm.
- A cancellation fee of \$ 150.00 may be charged if the patient reschedules or cancels within five (5) business days, before the procedure appointment.

√ MEDICATIONS

- Do not take iron supplements for five (5) days prior to the procedure.
- Aspirin, diabetic medications/insulin, anticoagulation medications, heart medications, blood thinners; must be discussed with your Physician, Nurse Practitioner or Physician Assistant at the time of the pre-procedure office visit.
- Specified medications should be taken the morning of the procedure, as prescribed, **3 or more hours prior to the procedure start time with a small sip of water.**
- Do not stop any medications unless instructed.

√ 3 DAYS BEFORE THE PROCEDURE:

- Adjust your diet to prepare for the procedure. Stop eating high fiber foods including, seeds, quinoa, nuts, beans, corn/popcorn, raw fruits, raw vegetables and bran for **three (3) days prior** to the procedure.

√ THE DAY BEFORE THE PROCEDURE:

- This procedure requires that you follow the following diet(s) on the day before the procedure. Your doctor will determine which diet you should follow.

<input type="checkbox"/> Clear Liquid Diet	Breakfast	Lunch	Dinner
<input type="checkbox"/> Low Residue Diet	Breakfast	Lunch	Dinner – before 6 pm

ALLOWED ON A CLEAR LIQUID DIET: apple juice, white grape and white cranberry juice, broth, tea, coffee without milk, creamer, Jell-O, Italian Ices, popsicles, sodas, Kool-Aid, Gatorade, and water.
You may continue these clear liquids up until 4 hours prior to your procedure.

NOT ALLOWED ON A CLEAR LIQUID DIET: red, orange, and purple colored liquids, as well as the following products, milk, cream, soy milk equivalent, artificial creamer, juices containing pulp (i.e. orange, grapefruit, pineapple, tomato & V-8 juice). Solid foods are not allowed on a clear liquid diet.

ALLOWED ON A LOW RESIDUE DIET: Ensure, Boost, eggs and plain or flavored yogurt without seeds or nuts. NO other foods are allowed on this diet.

SUPREP PREP INSTRUCTIONS – DO NOT FOLLOW THE INSTRUCTIONS ON THE BOX

√ **FOLLOW THE TABLE BELOW THE DAY PRIOR TO THE PROCEDURE:**

First Dose 6:00 pm the day before the Procedure	Pour the entire contents of one (1) bottle of Suprep (6oz) bowel prep into the mixing container provided. Fill the container with water or any clear liquid listed (10oz) to the 16-oz. fill line. Drink the entire amount over the next 30 minutes.
	Drink at least two (2) 16 oz. glasses of water or clear liquid over the next 2 hours.

√ **FOLLOW THE TABLE BELOW SIX (6) HOURS PRIOR TO THE PROCEDURE:**

Second Dose Six (6) hours prior to your scheduled procedure time _____	Pour the entire contents of one (1) bottle of Suprep (6oz) bowel prep into the mixing container provided. Fill the container with water or any clear liquid listed (10oz) to the 16-oz. fill line. Drink the entire amount over the next 30 minutes.
Four (4) hours prior to your scheduled procedure time _____	Drink at least two (2) 16 oz. glasses of water or clear liquid. You must be finished drinking 4 hours prior to your scheduled procedure time.
Additional Instructions:	

√ **WHAT TO EXPECT:**

- Expect to have frequent bowel movements and diarrhea within 1-3 hours. Be patient and try walking around to stimulate bowel motility. Remain near a restroom.
- Expect that some patients will have a delayed response. If you have not moved your bowels within 3 hours after taking the first dose then call the office.
- Expect to pass clear or yellow fluid at the completion of your prep.
- You must also take the second dose of the prep even if you are passing clear or yellow fluid.
- Expect that patients who do not fully complete their prep and are not adequately prepared may be canceled or rescheduled.

√ **WHAT TO DO TO MAKE PREPPING EASIER:**

- Patients can use Desitin, Anusol, Tucks pads or Vaseline to coat the rectal area to avoid irritation during the prep.
- Patients can refrigerate the prep to make drinking the solution easier.
- Patients can mix the solution with Sprite, Ginger ale or Lemonade.
- Patients can drink the solution with a straw or hold your nose when drinking.
- Patients should slow down and take time between doses if feeling nauseous.
- Patients may continue drinking clear liquids up to four hours before the procedure. During the 4 hours before your procedure, take nothing by mouth, not even water, chewing gum or candy. **If you do, the procedure will be canceled.**

√ **LAST MINUTE CHECK LIST:**

- If a physician has prescribed for you an inhaler for asthma, bring it with you to the procedure.
- **Take your blood pressure, heart, Parkinson’s, seizure or asthma medications, please take these 3 or more hours prior to the procedure start time with a small sip of water.**
- Bring with you your photo ID, insurance cards, co-payment if required.
- Leave valuables and jewelry at home and wear comfortable clothing.
- Avoid wearing perfume or lotions on the day of your procedure.

PLEASE LIST:

NAME OF ESCORT HOME: _____ **RELATIONSHIP:** _____

PHONE NUMBER: _____