

Jeffrey Garbis, M.D. Jeffrey Bernstein, M.D. Marvin Lawrence, II, M.D. Priti Bijpuria, M.D. Darryn Potosky, M.D. Richard Chasen, M.D. Theodore Kim, M.D. Sean Karp, M.D. Lauren Halvorson, M.D. Eileen Erskine, C.R.N.P.

Colonoscopy with OSMOPREP EGD & Colonoscopy with OSMOPREP

Patient Name:				
Date:	Arrival Time fo	Arrival Time for Procedure:		
With Doctor:				
Location:	Ambulatory Endoscopy Center of Maryland	Laurel Regional Hospital		
	Washington Adventist Hospital	Howard County General		

YOU MUST FILL YOUR PRESCRIPTION FOR OSMOPREP

You will be given sedation for your procedure and <u>MUST</u> have a ride <u>FROM</u> the examination. You <u>CANNOT</u> drive yourself home and <u>MUST</u> be accompanied by a responsible adult. You <u>MAY NOT</u> go home on public transportation alone. Pick up time will be approximately 1 hour after the procedure start time.

In order to be properly prepared for your procedure, please review the Pre-Procedure requirements on the attached sheet <u>one week</u> prior to your exam and follow the instructions carefully.

The liquids in the prep will cause you to have repeated loose stools (diarrhea), which is the expected result. These solutions typically work within 30-60 minutes but may take as long as 4 hours to have an effect. During the preparation process, you are advised to have a bathroom nearby. <u>It is very important that you follow each step and complete all of these instructions or the colonoscopy may have to be repeated.</u> The bowel preparation will not work properly unless you <u>drink plenty of liquids.</u>

Should you have any questions please contact our office at 301-498-5500.

If you should need to reschedule your procedure, we require at least 48 hour notice. If you fail to report on your procedure day or if you cancel your procedure with less than 48 hour notice, you may be subject to a \$150.00 late cancellation fee.

<u>Address & Directions to Ambulatory Endoscopy Center of MD:</u> **7350 Van Dusen Road, Suite 230,** Laurel, MD 20707. Take I 95 to exit 33 A, Route 198 towards Laurel. At first intersection, turn right onto Van Dusen Road. At 5th traffic light, make a right into the Laurel Regional Hospital driveway. The Medical Arts Pavilion is on the right. You should report to Suite 230.

COLONOSCOPY OSMOPREP CHECKLIST

CAPITAL DIGESTIVE CARE, LLC

Appointment Day: M Tu

Capital Digestive Care Pathology (pathology).

11/08/17 GG/RJ

W

F

Th

	Date: Arrival Tim	e:		
<u>Step 1:</u> (5) Five days before your colonoscopy: Date:				
 Please rea Fill prescr Discontin Discontin Inflammato You may to Discontin (e.g. Plavix, physician, b 	ad through entire instruction sheet iption for Osmoprep ue medications containing iron ue Ibuprofen, Advil/Motrin/Aleve & Anti- ry Medications & other NSAIDS use Tylenol (Acetaminophen) for discomfort ue blood thinners/anticoagulation medications 'Coumadin/Warfarin/Eliquis) as instructed by your out take all other medications including blood	Step 2: (3) Three days before colonoscopy: Date: □ Stop eating high fiber foods including: nuts, corn, popcorn, raw fruits, raw vegetables & bran Notes:		
□ Notes:				
(1) One da (1) One da No solid fu In the mo blue or purp Examples products), c carbonated Dew (regula fruit juices v (yellow, gre Drink 6-8 dehydratior At 6:00 pu regimen dia liquid every At 7:00 pu 40 oz. of cle Unit bedtim	 <u>n</u>: Start the first dosing regimen (See first dosing gram). Take 4 tablets with at least 8 oz. of clear 15 minutes for 1 hour <u>n</u>: You should have taken a total of 20 tablets and 	Step 4: Day of your colonoscopy: Date: NO gum, mints, hard candy or smoking day of procedure At am/pm (5 hours before your procedure): Take 4 tablets with at least 8 oz. of clear liquids every 15 minutes for 30 minutes, then STOP drinking clear liquids. (See second dosing regimen diagram) After 30 minutes, you should have taken a total of 12 tablets and 24 oz. of clear liquid. After finishing your second dosing regimen, take all other medications including blood pressure medication with sips of water DO NOT eat, smoke or chew anything by mouth 8 hours before your procedure You may drink clear liquids up to 4 hours before leaving for your procedure. Stop time:		
	e a total of 4 statements; (1) Ambulatory Endoscopy Center of MD bital Digestive Care (physician), (3) Corridor Anesthesia (anesthesia) & (4)	Notes:		