#### **Clear Liquid Diet**

You may have any of the following the day before your colonoscopy:

- Regular or Diet Ginger Ale, Sprite or 7-Up (no cherry flavor)
- Water
- Apple juice
- ☐ Gatorade (not orange, red or purple)
- □ Popsicles (not orange, red or purple)
- ☐ Chicken or vegetable consommé or clear broth
- Yellow or green Jell-O
- Tea
- White grape juice
- □ Lemonade (not pink lemonade and no pulp)

## Preparing for your Colonoscopy Helpful Hints....

- Remember the goal of the prep is to get cleaned out. When the color of what you are passing is clear or cloudy yellow with some flecks, then you are clean. If you are passing brown stool or brown water then you may not be adequately prepared. However, you will still need to do the second dosing as directed on the other side even if you are clear the night before.
- Sucking on a lemon drop or other hard candy or a fresh lemon or lime between glasses of the solution can help clear any aftertaste.
- Nausea may occur when multiple glasses of the solution have been taken and nothing has come out rectally. Walking will help move the solution through the body.
- It is helpful to put Vaseline around the anal area to minimize irritation from frequent bowel movements.
- If you have any problems with the bowel cleansing process during the daytime, you can contact the office. If you have difficulty in the evening, please contact the physician "on-call" by calling the after hours number 1-877-722-7098.

#### **Colonoscopy Checklist**

There is additional information in your colonoscopy instruction packet.

For other questions or to schedule an office visit, call

240-737-0085

Detailed information is also available online at: www.CCendo.com or www.Endodc.com



First in Digestive Health

#### Metropolitan Gastroenterology Group

- A Division of Capital Digestive Care, LLC -

## Your Five (5) Day Colonoscopy Checklist

# OSMO Preparation Instructions



Chevy Chase Endoscopy Center

The Endoscopy Center of Washington DC

Bethesda Endoscopy Center

Sibley Hospital

George Washington Hospital

Suburban Hospital

### Your Five (5) Day Colonoscopy Checklist

Continue your medications unless instructed other	
Continue your medications unless instructed ou	nerwise by your Physician.
Three (3) days before your Colonoscopy  Stop fiber supplements such as Metamucil, Fiber Purchase your OSMO pills (prescription enclose)	
The day before your Colonoscopy	
turkey sandwich on white bread (bread and magnetic After 12 noon – CLEAR LIQUIDS ONLY (see list of the second seco	eam substitutes – NO DAIRY PRODUCTS. on – NO LUNCH. , you may have the following for lunch: small chicken or neat only). of clear liquids on back of this page). will be taking 20 tablets. Remember to drink 8 oz of
□ 6 pm – Take 4 pills w/8 oz of liquid □ 6:30 pm – Take 4 pills w/8 oz of liquid □ 7:00 pm – Take 4 pills w/8 oz of liquid	<ul> <li>□ 6:15 pm – Take 4 pills w/8 oz of liquid</li> <li>□ 6:45 pm – Take 4 pills w/8 oz of liquid</li> </ul>
	arrhea this evening. Some individuals will have a delayed the medication, be patient and try walking around to stimulate ellow fluid tonight, you must take the rest of your prep tomorrow.
The day of your Colonoscopy	
<ul> <li>□ Remember, take your medications unless your process.</li> <li>□ 4 hours before your arrival time, you will take the Remember to drink 8 oz of apple juice, ginger</li> <li>□ 1st dose – Take 4 pills w/8 oz of liquid</li> </ul>	ne remaining 12 pills, 4 pills every 15 minutes.  r ale, water or Gatorade with each dose
<ul> <li>Remember, take your medications unless your personant description.</li> <li>4 hours before your arrival time, you will take the Remember to drink 8 oz of apple juice, ginger</li> </ul>	ne remaining 12 pills, 4 pills every 15 minutes. r ale, water or Gatorade with each dose w/8 oz of liquid
<ul> <li>Remember, take your medications unless your presented 4 hours before your arrival time, you will take the Remember to drink 8 oz of apple juice, ginger</li> <li>□ 1st dose - Take 4 pills w/8 oz of liquid</li> <li>□ 2nd dose - 15 min later - Take 4 pills</li> <li>□ 3rd dose 15 min after 2nd dose - Take</li> </ul>	me remaining 12 pills, 4 pills every 15 minutes.  r ale, water or Gatorade with each dose  w/8 oz of liquid e 4 pills w/8 oz of liquid
<ul> <li>Remember, take your medications unless your presented 4 hours before your arrival time, you will take the Remember to drink 8 oz of apple juice, ginger</li> <li>□ 1st dose - Take 4 pills w/8 oz of liquid</li> <li>□ 2nd dose - 15 min later - Take 4 pills</li> <li>□ 3rd dose 15 min after 2nd dose - Take</li> </ul>	me remaining 12 pills, 4 pills every 15 minutes.  r ale, water or Gatorade with each dose  w/8 oz of liquid e 4 pills w/8 oz of liquid
Remember, take your medications unless your processed 4 hours before your arrival time, you will take the Remember to drink 8 oz of apple juice, ginger  1st dose - Take 4 pills w/8 oz of liquid 2nd dose - 15 min later - Take 4 pills 3rd dose 15 min after 2nd dose - Take Two-and-a-half (2-1/2) hours before your ARF  DO NOT DRINK OR EAT ANYTHING	me remaining 12 pills, 4 pills every 15 minutes.  r ale, water or Gatorade with each dose  w/8 oz of liquid e 4 pills w/8 oz of liquid
Remember, take your medications unless your processed 4 hours before your arrival time, you will take the Remember to drink 8 oz of apple juice, ginger  1st dose - Take 4 pills w/8 oz of liquid 2nd dose - 15 min later - Take 4 pills 3rd dose 15 min after 2nd dose - Take Two-and-a-half (2-1/2) hours before your ARF  DO NOT DRINK OR EAT ANYTHING	me remaining 12 pills, 4 pills every 15 minutes.  r ale, water or Gatorade with each dose  w/8 oz of liquid e 4 pills w/8 oz of liquid  RIVAL TIME
Remember, take your medications unless your partial time, you will take the Remember to drink 8 oz of apple juice, ginger  1st dose – Take 4 pills w/8 oz of liquid 2nd dose – 15 min later – Take 4 pills 3rd dose 15 min after 2nd dose – Take Two-and-a-half (2-1/2) hours before your ARF DO NOT DRINK OR EAT ANYTHING  Last Minute Checklist  Bring this checklist with the name and phone numbers	me remaining 12 pills, 4 pills every 15 minutes.  r ale, water or Gatorade with each dose  w/8 oz of liquid e 4 pills w/8 oz of liquid  RIVAL TIME
Remember, take your medications unless your processed to the second seco	me remaining 12 pills, 4 pills every 15 minutes.  r ale, water or Gatorade with each dose  w/8 oz of liquid e 4 pills w/8 oz of liquid  RIVAL TIME
Remember, take your medications unless your processed to the second seco	me remaining 12 pills, 4 pills every 15 minutes.  r ale, water or Gatorade with each dose  w/8 oz of liquid e 4 pills w/8 oz of liquid  RIVAL TIME  Phone Number:  (husband, wife, sibling, friend, significant other, child, etc)  ayment required for the procedure.  medications you are currently taking.

Patient's signature